
Complementarity in LLP Searches

Marco Drewes, Université catholique de Louvain Snowmass EF09 Meeting August 7 2020

Overview

- Portal Effective Theory
- Constraints on LLPs from Cosmology
- Global Constraints
- LLPs in Heavy Ion Collisions

Portal Effective Theory



- many scenarios predict comparably light "Hidden Sectors" that give rise to Long Lived Particles (LLPs)
- Hidden Sector couples to SM via "portals"
- A single "portal" operator at high energies can give rise to different interactions at low energies

Desirable: A general EFT framework that maps fundamental parameters at high energies onto low energy observables

Portal Effective Theory

Electroweak Scale PETs
$$d \leq 5$$

 \Downarrow

General GeV Scale PETs
$$(d \le 7 + NDA)$$


 \Downarrow

$$\left\{ \begin{array}{l} \mathsf{Specific\ PETs} \\ (\chi\mathsf{PT,\ HQET,\ etc.}) \end{array} \right\}$$

- \blacksquare Suppressed by small parameter ϵ_{UV}
- d ≤ 5 captures axion-like particles (ALPs), heavy neutral leptons (HNLs), dark photons, etc.
- d = 6,7 Operators capture $1/M_W^2$ suppressed $\Delta F \neq 0$ interactions
- Naive Dimensional Analysis (NDA) counts (4π) loop suppression factors (keep only leading order)
- QCD nonperturbative ⇒ Specific PETs
- \blacksquare PET χ PT captures meson interactions at NA62 in Kaon mode


LLPs in Cosmology

- LLPs can affect cosmological observables through their presence (expansion rate of the universe) or decays (entropy injection, disturbance of distribution function, dissociation of nuclei...)
- Observables: Neff and light element abundances in intergalactic medium
- This can provide a "floor" for experimental searches
- CMB S4 experiment will set new bounds




...combined BBN + CMB studies planned with Isabel Oldengott

Global Constraints



Combined observables from different sources can

- identify full exclusion potential of future experiments by projecting improvement in direct and indirect searches
- In case of a discovery: Can help to pin down underlying model (minimal type-I seesaw is fully testable)
- Example: RH neutrino mass and mixing




...further studies planned with M. Chrzaszcz, T. Gonzalo, J. Harz., C. Weniger

LLPs in Heavy Ion Collisions

Heavy Ion collisions have a number of advantages: Bruce et al 1812.07688

- new production mechanisms (e.g. in strong fields)
- different backgrounds (no pile-up, ...)
- very low triggers



...ongoing studies with Jan Hajer, Andrea Giammanco, Hesham el Faham

dedicated workshop at ECT* May 17-21, 2021 https://indico.cern.ch/event/831940/